

RESEARCH ARTICLE

ANTIFUNGAL POTENTIAL OF EXO-METABOLITES PRODUCED BY *PENICILLIUM CAPSULATUM* AGAINST PHYTOPATHOGENIC FUNGI

Niharika Mohanty, Atmaja Elina Mishra and Nibha Gupta*

Plant Pathology and Microbiology Division, Regional Plant Resource Centre, Bhubaneswar, Odisha-751015

*Corresponding author: nguc2003@yahoo.co.in

This is an open access journal distributed under the Creative Commons Attribution License CC BY 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

ARTICLE DETAILS

Article History:

Received 12 August 2024
Revised 17 September 2024
Accepted 19 October 2024
Available online 06 November 2024

ABSTRACT

Fungi are well-known producers of secondary metabolites, commonly referred to as exo-metabolites, which play crucial roles in ecological interactions, including plant protection. Among these metabolites, many exhibit potent antifungal properties, making them effective biocontrol agents against phytopathogenic fungi that cause significant agricultural losses. This study investigates the antifungal potential of solvent extracts derived from the extracellular secondary metabolites of *Penicillium capsulatum*, specifically targeting five phytopathogenic fungi: *Fusarium equseti*, *Fusarium acuminatum*, *Colletotrichum gloeosporioides*, *Fusarium javanicum*, and *Nigrospora sphaeria*. Using a solvent extraction method, partially purified fungal metabolites were tested for their antifungal activity via the pour plate method. The extracts were incorporated into agar medium in two concentrations: 100µl (lower dose) and 500µl (higher dose), and the growth reduction percentage was calculated based on the colony diameter measurements. The results indicated that the toluene extract of *P. capsulatum* exhibited the highest growth reduction against *F. equseti*, achieving $45.00 \pm 3.54\%$ at the higher dose and $37.50 \pm 3.54\%$ at the lower dose. The methanolic extract demonstrated complete inhibition ($100.00 \pm 0.00\%$) of *F. acuminatum* and *C. gloeosporioides* at the higher dose, while acetic acid, DMSO, and acetonitrile extracts showed varying levels of antifungal activity against other target pathogens. These findings highlight the potential of *P. capsulatum* metabolites as effective antifungal agents against diverse phytopathogenic fungi, presenting a promising alternative to chemical fungicides. Furthermore, the results support the need for further exploration of fungal metabolites in the development of sustainable biocontrol strategies for managing plant diseases.

KEYWORDS

Fungi, Antagonism, Metabolite, *Penicillium capsulatum*, Growth reduction.

1. INTRODUCTION

Fungi produce a wide range of secondary metabolites, commonly referred to as exo-metabolites, which play significant roles in their ecological interactions. Among these metabolites, many possess antifungal properties that can be particularly effective against phytopathogenic fungi i.e. organisms that cause diseases in plants (Chen et al., 2021). Phytopathogenic fungi are responsible for substantial agricultural losses worldwide, as they lead to plant diseases like rusts, blights, and wilts, which severely impact crop yields (Mahadevakumar and Sridhar 2021).

Exo-metabolites, including enzymes, organic acids, and antimicrobial compounds, can inhibit the growth of these harmful fungi, making them valuable tools in biocontrol strategies. These metabolites often interfere with the cell walls or metabolic processes of the target fungi, leading to their suppression or death (Bharadwaj et al., 2021). Utilizing such naturally derived antifungal compounds offers a more sustainable and environmentally friendly alternative to synthetic chemical fungicides, which are known to have adverse effects on ecosystems and can lead to resistance in pathogenic strains (Jimenez-Reyes et al., 2019).

Numerous endophytic fungi have been identified for their potent antifungal properties. For instance, ergot alkaloids produced by the fungus *Claviceps purpurea* have been found to exhibit strong antifungal activity (Dembitsky 2014). Additionally, acetylenic metabolites produced by *Penicillium* and *Aspergillus* species, along with other filamentous fungi, have emerged as a distinct chemical family of antifungal compounds.

These metabolites have even inspired the development of synthetic fungicides due to their unique structure and bioactivity (Baldt et al., 2013; Kuklev et al., 2013). In recent years, new discoveries continue to highlight the potential of fungi as biocontrol agents. For example, capsulactone, isolated from the endophytic fungus *Penicillium capsulatum* XL027, which was obtained from the leaves of *Panax notoginseng*, exhibited antimicrobial activity against methicillin-resistant *Staphylococcus aureus*. Although its antimicrobial activity was modest, capsulactone represents another important example of the capacity of biocontrol fungi to produce bioactive secondary metabolites with potential applications in combating pathogens (Xian et al., 2021).

The ability of fungi to produce bioactive secondary metabolites has long been recognized, with griseofulvin being one of the earliest reported examples. This compound, classified as a benzofuran, was isolated from *Penicillium* species and demonstrated antifungal activity by inhibiting the growth of the plant pathogen *Botrytis cinerea*, leading to abnormal hyphal growth and disorientation (Rodrigo et al., 2021). This discovery significantly advanced the understanding of fungal secondary metabolites and their role as biocontrol agents against phytopathogens. These findings underscore the chemical diversity of fungal metabolites and their continued relevance as a resource for developing new biocontrol agents to manage both plant and microbial pathogens (Rush et al., 2021).

Thus, the exploration of exo-metabolites produced by fungi as antifungal agents holds promise in developing novel plant protection methods, which can mitigate the threats posed by phytopathogenic fungi while promoting

Quick Response Code



Access this article online

Website:
www.actascientificamalaysia.com

DOI:
10.26480/asm.02.2024.67.70

sustainable agricultural practices.

2. MATERIALS AND METHODS

2.1 Microorganisms

This experiment was conducted by selecting a biocontrol fungi *Penicillium capsulatum* and tested against five phytopathogenic fungi: *Fusarium equiseti*, *Fusarium acuminatum*, *Colletotrichum gloeosporioides*, *Fusarium javanicum*, and *Nigrospora sphaeria*. These were collected from Microbiology and Plant Pathology Laboratory of Regional Plant Resource Centre, Bhubaneswar, Odisha.

2.2 Solvent Extraction Method

Extracellular secondary metabolites of *Penicillium capsulatum* were separated through solvent extraction and partially purification by column chromatography. Mass scale production of the fungal strain *Penicillium capsulatum* was done. This organism was inoculated in Sabouraud dextrose broth medium and incubated at 30°C. After incubation period, filtration and solvent extraction was done. Partial purification was done through column chromatography process by passing the crude extract using polar to nonpolar solvents (Acetic acid, methanol, ethanol, isopropanol, acetonitrile, acetone, chloroform, pyridine, DMSO, ethyl acetate, toluene and hexane) (Chapla et al., 2014). Prepared extracts were tested for antifungal activity against five numbers of phytopathogenic fungi by following pour plate method. Plates were prepared by adding different solvent extracts in molten agar medium in two doses i.e. 100µl (lower dose) and 500µl (higher dose). 6mm disc cuttings of the well grown fungal culture were placed at the middle of the plate. Colony diameter of the organisms was measured and percentage of growth reduction was calculated (Gu et al., 2022).

3. RESULTS AND DISCUSSION

This study demonstrates the significant antifungal potential of various

solvent extracts from *Penicillium capsulatum* against five key phytopathogenic fungi, including *Fusarium equiseti*, *Fusarium acuminatum*, *Colletotrichum gloeosporioides*, *Fusarium javanicum*, and *Nigrospora sphaeria*. The results reveal that among the extracts tested, the toluene extract exhibited the highest growth inhibition against *F. equiseti*, with a reduction percentage of $45.00 \pm 3.54\%$ at a higher dose and $37.50 \pm 3.54\%$ at a lower dose. This suggests that toluene extract may contain bioactive compounds that are highly effective against this particular pathogen. Interestingly, the chloroform extract showed no inhibitory effect at a lower dose but exhibited some activity at the higher dose, albeit with a lower reduction percentage of $14.03 \pm 0.00\%$ (figure 1). In contrast, *F. acuminatum* was completely inhibited by the methanolic extract at a higher dose ($100.00 \pm 0.00\%$), followed by notable inhibition from the acetic acid extract. Toluene ($55.00 \pm 1.77\%$) and DMSO ($48.75 \pm 1.77\%$) extracts also displayed moderate antifungal activity at lower doses (figure 2). These results indicate the versatility of *P. capsulatum* extracts, particularly methanol which effectively inhibited multiple phytopathogens, including *Colletotrichum gloeosporioides* up to 100.00%, and *F. javanicum*, where the methanolic extract demonstrated high efficacy at both lower and higher doses (figure 3 and 4). Additionally, acetonitrile, acetone, and ethyl acetate extracts also exhibited antifungal activity, particularly against *F. javanicum* and *N. sphaeria*. Notably, 100% growth inhibition of *N. sphaeria* was observed with acetic acid extract at both dose levels, further highlighting its potential as a potent biocontrol agent (figure 5).

Complementing these findings, previous research has shown that ethyl acetate extracts from various fungal species exhibit similar biocontrol properties against a range of phytopathogens. For example, pretrichoderamide A, isolated from *Trichoderma harzianum* and *Epicoccum nigrum*, has demonstrated antimicrobial efficacy against *Ustilago maydis*, a common plant pathogen (Harwoko et al., 2019). Similarly, ethyl acetate extracts from *Fusarium oxysporum* endophytes have been shown to contain palmitic, stearic, and octadecenoic acids, which are known to act as potent fungicides (Killic et al., 2021).




Figure 1: Effect of solvent extracts of *P. capsulatum* against *Fusarium equiseti*




Figure 2: Effect of solvent extracts of *P. capsulatum* against *Fusarium acuminatum*




Figure 3: Effect of solvent extracts of *P. capsulatum* against *Colletotrichum gloeosporioides*




Figure 4: Effect of solvent extracts of *P. capsulatum* against *Fusarium javanicum*




Figure 5: Effect of solvent extracts of *P. capsulatum* against *Nigrospora spherica*

These fatty acids, along with other metabolites such as leucinostatin produced by *Purpureocillium lilacinum*, exhibit broad-spectrum activity against both bacterial and fungal pathogens, further highlighting the role of fungal metabolites in plant protection (Wang et al., 2016). Moreover, *Metarhizium anisopliae* has emerged as an effective biocontrol agent, particularly against *Fusarium graminearum*, due to its ability to produce fungistatic secondary metabolites, enhance wheat growth, and trigger defense responses in plants (Hao et al., 2021). Additionally, the susceptibility of other plant pathogens such as *Alternaria solani*,

Rhizoctonia solani, *Pythium ultimum*, and *Colletotrichum lagenarium* to fatty acids produced by fungal endophytes emphasizes the broad biocontrol potential of fungal metabolites (Raynor et al., 2004; Liu et al., 2008).

In summary, the findings from this study, coupled with the existing body of research, underscore the substantial role of fungal secondary metabolites as effective biocontrol agents. The use of solvent extracts, particularly those derived from *P. capsulatum* and other fungi, offers a

promising alternative to chemical fungicides for managing plant diseases.

4. CONCLUSION

The findings from this study highlight the considerable antifungal potential of *Penicillium capsulatum* solvent extracts against a range of phytopathogenic fungi, including *Fusarium equseti*, *Fusarium acuminatum*, *Colletotrichum gloeosporioides*, *Fusarium javanicum*, and *Nigrospora sphaeria*. The highest level of inhibition was observed with the toluene and methanolic extracts, particularly in the case of *F. equseti* and *F. acuminatum*, where methanol demonstrated complete growth inhibition at a higher dose. The versatility of *P. capsulatum* extracts across different solvents, particularly methanol, acetic acid, and toluene, underscores the effectiveness of these natural compounds in managing various plant pathogens. The complete inhibition of *N. sphaeria* by acetic acid further reinforces the potential of these fungal secondary metabolites as potent biocontrol agents. These results demonstrate the value of exploring fungal exo-metabolites as an eco-friendly alternative to chemical fungicides, offering an avenue towards sustainable agricultural practices. The efficacy of these natural antifungal agents provides a promising strategy to reduce the harmful environmental impacts associated with synthetic fungicides while ensuring effective pathogen control. Continued research is needed to further optimize these extracts for practical applications, as well as to investigate their modes of action and broader spectrum of activity, contributing to innovative and sustainable plant protection solutions.

ACKNOWLEDGMENT

The authors are grateful to Forest and Environment Department, Govt. of Odisha for financial assessment through State Plan project and the Chief Executive, RPRC for providing laboratory and administrative facilities.

REFERENCES

Azerang, P., and Sardari, S. 2012. Antifungal activity of enynediesters and acetylenic compounds obtained by synthesis and in silico prediction pattern. *J. Mycol. Med.*, 22, Pp. 230–236. <https://doi.org/10.1016/j.mycmed.2012.06.001>

Bharadwaj, R., Kumar, S.R., Sathishkumar R. 2021. Insights into Role of Invisible Partners in Plant Growth and Development. *Plant Growth and Stress Physiology*, Pp. 23-48. https://doi.org/10.1007/978-3-030-78420-1_2

Bladt, T.T., Frisvad, J.C., Knudsen, P.B., Larsen, T.O. 2013. Anticancer and antifungal compounds from *Aspergillus*, *Penicillium* and other filamentous fungi. *Molecules*, 18, Pp. 11338–11376. <https://doi.org/10.3390/molecules180911338>

Chapla, V.M., Zeraik, M.L., Leptokarydis, I.H., Silva, G.H., Bolzani, V.S., Young, M.C., Pfenning, L.H., Araújo, A.R. 2014. Antifungal compounds produced by *Colletotrichum gloeosporioides*, an endophytic fungus from *Michelia champaca*. *Molecules*, 19(11), Pp. 19243–52. <https://doi.org/10.3390/molecules191119243>

Chen, L., Schwier, M., Krumbach, J., Kopriva, S., Jacoby, R.P. 2021. Metabolomics in plant-microbe interactions in the roots. *Advances in Botanical Research*, 98, Pp. 133–61. <https://doi.org/10.1016/bs.abr.2020.09.018>

Dembitsky, V.M. 2014. Naturally occurring bioactive Cyclobutane-containing (CBC) alkaloids in fungi, fungal endophytes, and plants. *Phytomedicine*, 21, Pp. 1559–1581. <https://doi.org/10.1016/j.phymed.2014.07.005>

Gu, H., Zhang, S., Liu, L., Yang, Z., Zhao, F., Tian, Y. 2022. Antimicrobial potential of endophytic fungi from *Artemisia argyi* and bioactive metabolites from *Diaporthe* sp. AC1. *Frontiers in Microbiology*, 13, Pp. 908836. <https://doi.org/10.3389/fmicb.2022.908836>

Hao, Q., Albaghdady, D.M.D., Xiao, Y., Xiao, X., Mo, C., Tian, T., Wang, G. 2021. Endophytic *Metarhizium anisopliae* is a potential biocontrol agent against wheat *Fusarium head blight* caused by *Fusarium graminearum*. *J. Plant Pathol.* 103, Pp. 875–885. <https://doi.org/10.1007/s42161-021-00866-6>

Harwoko, H., Daletos, G., Stuhldreier, F., Lee, J., Wesselborg, S., Feldbrügge, M., Müller, W.E.G., Kalscheuer, R., Ancheeva, E., Proksch, P. 2019. Dithiodiketopiperazine derivatives from endophytic fungi *Trichoderma harzianum* and *Epicoccum nigrum*. *Nat. Prod. Res.* 35, Pp. 257–265. <https://doi.org/10.1080/14786419.2019.1627348>

Jimenez-Reyes, M.F., Carrasco, H., Olea, A.F., Silva-Moreno, E. 2019. Natural compounds: A sustainable alternative to the phytopathogens control. *Journal of the Chilean Chemical Society*, 64(2), Pp. 4459–65. <https://doi.org/10.4067/s0717-97072019000204459>

Kilic, G., Tosun, G., Bozdeveci, A., Erik, I., Ozturk, E., Reis, R., Sipahi, H., Cora, M., Karaoglu, S.A., Yayli, N. 2021. Antimicrobial, cytotoxic, antiviral effects, and spectroscopic characterization of metabolites produced by *Fusarium oxysporum* YP9B. *Rec. Nat. Prod.*, 15, Pp. 547–567. <https://doi.org/10.25135/rnp.208.20.06.1674>

Kuklev D.V., Domb, A.J., Dembitsky, V.M. 2013. Bioactive acetylenic metabolites. *Phytomedicine*, 20, Pp. 1145–1159. <https://doi.org/10.1016/j.phymed.2013.06.009>

Liu, S., Ruan, W., Li, J., Xu, H., Wang, J., Gao, Y., Wang, J. 2008. Biological control of phytopathogenic fungi by fatty acids. *Mycopathologia* 166, Pp. 93–102. <https://doi.org/10.1007/s11046-008-9124-1>

Mahadevakumar, S., and Sridhar, K.R. 2021. Diversity of Pathogenic Fungi in Agricultural Crops. *Plant, Soil and Microbes in Tropical Ecosystems*, Pp. 101-49. https://doi.org/10.1007/978-981-16-3364-5_6

Rodrigo, S., Garcia-Latorre, C., Santamaria, O. 2021. Metabolites produced by fungi against fungal phytopathogens: Review, implementation and perspectives. *Plants*, 11(1), Pp. 81. <https://doi.org/10.3390/plants11010081>

Rush, T.A., Shrestha, H.K., Gopalakrishnan, M.M., Spangler, M.K., Ellis, J.C., Labbe, J.L., Abraham, P.E. 2021. Bioprospecting *Trichoderma*: A systematic roadmap to screen genomes and natural products for biocontrol applications. *Frontiers in Fungal Biology*, 716511. <https://doi.org/10.3389/ffunb.2021.716511>

Raynor, L., Mitchell, A., Walker, R., Walker, K. 2004. Antifungal activities of four fatty acids against plant pathogenic fungi. *Mycopathologia*, 157, Pp. 87–90. <https://doi.org/10.1023/b:myco.0000012222.68156.2c>

Wang, G., Liu, Z., Lin, R., Li, E., Mao, Z., Ling, J., Yang, Y., Yin, W.B., Xie, B. 2016. Biosynthesis of antibiotic leucinoastatins in bio-control fungus *Purpureocillium lilacinum* and their inhibition on *Phytophthora* revealed by genome mining. *PLoS Pathog.*, 12, Pp. e1005685. <https://doi.org/10.1371/journal.ppat.1005685>

Xian, P.J., Chen, H.Y., Feng, Z., Zhao, W., Yang, X.L. 2021. Capsulactone: a new 4-hydroxy-a-pyrone derivative from an endophytic fungus *Penicillium capsulatum* and its antimicrobial activity. *Journal of Asian Natural Products Research*, 23(11), Pp. 1100–1106. <https://doi.org/10.1080/10286020.2020.1847092>

Xu, T., Tripathi, S.K., Feng, Q., Lorenz, M.C., Wright, M.A., Jacob, M.R., Mask, M.M., Baerson, S.R., Li, X.C., Clark, A.M. 2012. A potent plant-derived antifungal acetylenic acid mediates its activity by interfering with fatty acid homeostasis. *Antimicrob. Agents Chemother.*, 56, Pp. 2894–2907. <https://doi.org/10.1128/aac.05663-11>

